

OXERVATE® is a prescription eye drop solution used to treat a condition called neurotrophic keratitis. OXERVATE is safe and effective in children two years of age and older.



Take a closer look at OXERVATE®

This brochure will help you understand why your doctor prescribed OXERVATE and how to properly use it

Not an actual patient.

Important Safety Information

What is the most important information I should know about OXERVATE?

- If you wear contact lenses in your affected eye or both eyes, remove them before using OXERVATE and **wait 15 minutes after** using OXERVATE before reinserting them.

Please see continuation of Important Safety Information on [page 2](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information for OXERVATE](#).

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

Disease
Information

About

Using

How to Use

Getting
& Storing

Important Safety
Information

Discussion
Guide

What to know about your treatment with OXERVATE®



Get help accessing OXERVATE®:

Dompé CONNECT to Care is a patient support program that helps people start treatment with OXERVATE.



Delivered to you by a specialty pharmacy:

Accredo Health Group, Inc., a specialty pharmacy, coordinates the delivery of OXERVATE directly to you.



Eye drops are applied using your Delivery System Kit:

After reading this brochure, be sure to talk with your doctor about how to use OXERVATE.

Important Safety Information (cont'd)

What is the most important information I should know about OXERVATE? (cont'd)

- OXERVATE may cause an eye reaction such as eye pain. Talk to your doctor if you experience a more serious eye reaction.
- If you get an eye infection while using OXERVATE, talk to your doctor right away.

What should I avoid while using OXERVATE?

- Your vision may be blurred for a short time after using OXERVATE. If this happens, **wait until your vision clears before you drive or use machines.**

Please see continuation of Important Safety Information on [page 7](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

Table of contents

Disease Information	<u>4</u>
What Is Neurotrophic Keratitis (NK)?	<u>4</u>
About NK	<u>5</u>
About	<u>6</u>
What Is OXERVATE?	<u>6</u>
Clinical Trial Efficacy & Safety Information	<u>8</u>
Using	<u>10</u>
Using OXERVATE	<u>11</u>
Dosing Schedule	<u>13</u>
How to Use	<u>14</u>
Getting & Storing	<u>18</u>
Dompé CONNECT to Care & Accredo	<u>19</u>
The 5-Step Fulfillment Process	<u>20</u>
Storage	<u>21</u>
Important Safety Information	<u>23</u>
Doctor Discussion Guide	<u>24</u>

Please see continuation of Important Safety Information on [page 7](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.

oxervate® 
(cenegermin-bkbj ophthalmic
solution) 0.002% (20 mcg/mL)

NK and your cornea

Neurotrophic keratitis, or “NK,” is an eye disease that may be caused by damage to the corneal nerves. This damage may lead to poor healing and degeneration of the cornea. If not treated, it may lead to a corneal ulcer (an open sore in the cornea).

This image is for illustrative purposes only. This is not an exact representation of a patient.

NK is considered a rare disease and is estimated to affect approximately 70,000 people in the United States*

*Based on the IRIS® Registry (Intelligent Research in Sight) data published in 2022 and adapted number based on the US population.

Please see continuation of Important Safety Information on [page 7](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.


oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

Understanding your cornea & corneal nerves

The cornea is one of the most sensitive tissues in the body. As the outermost structure on the front of the eye, it protects the eye's inner contents.

Cornea

- Protects your eye from bacteria and dust
- Helps keep your eye from drying out
- Helps you see by focusing light into your eye



Corneal nerves

- Send signals to your brain to blink and produce tears to help maintain a healthy surface

NK may occur when the nerves of the cornea are damaged

When your corneal nerves are damaged, your cornea may have a lack of feeling and may be unable to send normal signals to your brain. This damage may lead to poor healing of the corneal surface. As a result, the corneal surface may break down, which may lead to NK.

Some people with NK may not complain of symptoms due to lack of feeling in the cornea.

Talk to your doctor about any symptoms you may be experiencing

Please see continuation of Important Safety Information on [page 7](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information for OXERVATE](#).

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)



What is OXERVATE®?

Please see continuation of Important Safety Information on [page 7](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

What is OXERVATE®?

The only prescription eye drop FDA-approved to treat neurotrophic keratitis (NK)



The active ingredient in OXERVATE® is cenegermin-bkbj

Cenegermin-bkbj is a manufactured form of a natural protein in the body called nerve growth factor (NGF).

If you have any questions about how to use OXERVATE, ask your doctor or pharmacist

Important Safety Information (cont'd)

What are the possible side effects of OXERVATE?

- The most common side effects of OXERVATE are eye pain, corneal deposits, feeling of having something in the eye, enlarged blood vessels in the white of the eyes, swelling of the eye, sensitivity to light, increase of tears, and headache.
- These are not all the possible side effects of OXERVATE.
- Tell your doctor if you have any side effects that bother you or do not go away.


Please see continuation of Important Safety Information on [page 9](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

OXERVATE® clinical trial results

In 2 clinical trials, 151 patients with neurotrophic keratitis (NK) were treated with OXERVATE® or placebo (inactive substance) for 8 weeks

Percentage of patients that experienced complete corneal healing with OXERVATE vs placebo at 8 weeks*



*At 8 weeks, each patient’s degree of corneal healing was determined by a test to assess damage to the cornea (corneal staining). If there was no staining in the damaged part of the cornea, and no persistent staining in the rest of the cornea, the cornea was considered completely healed (complete corneal healing).

More patients who received OXERVATE experienced complete corneal healing compared to those who received placebo (72% vs 33% and 65% vs 17%).

The most common side effect with OXERVATE was eye pain (~16%). Other side effects with OXERVATE (1% to 10%) included corneal deposits, feeling of having something in the eye, enlarged blood vessels in the white of the eyes, swelling of the eye, sensitivity to light, increase of tears, and headache.

Please see continuation of Important Safety Information on page 9 and complete Important Safety Information on page 23, as well as the full Prescribing Information, including Patient Information for OXERVATE.

oxervate® 
 (cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

OXERVATE side effects



This image is for illustrative purposes only. This is not an exact representation of a patient.

What are the possible side effects of OXERVATE?

- The most common side effects of OXERVATE are eye pain, corneal deposits, feeling of having something in the eye, enlarged blood vessels in the white of the eyes, swelling of the eye, sensitivity to light, increase of tears, and headache
- These are not all the possible side effects of OXERVATE
- Tell your doctor if you have any side effects that bother you or do not go away

Call your doctor for medical advice about side effects. You may report side effects to Dompé U.S. Inc at 1-833-366-7387 or FDA at 1-800-FDA-1088

General information about the safe and effective use of OXERVATE

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use OXERVATE for a condition for which it was not prescribed. Do not give OXERVATE to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or doctor for information about OXERVATE that is written for health professionals.

For more information about OXERVATE talk to your doctor or pharmacist.

The Doctor Discussion Guide on [page 24](#) may help.

Important Safety Information (cont'd)

What is the most important information I should know about OXERVATE?

- If you wear contact lenses in your affected eye or both eyes, remove them before using OXERVATE and **wait 15 minutes after** using OXERVATE before reinserting them.

Please see continuation of Important Safety Information on [page 10](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information for OXERVATE](#).

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)



Using OXERVATE®

Not an actual patient.

Important Safety Information (cont'd)

What is the most important information I should know about OXERVATE? (cont'd)

- OXERVATE may cause an eye reaction such as eye pain. Talk to your doctor if you experience a more serious eye reaction.
- If you get an eye infection while using OXERVATE, talk to your doctor right away.

Please see continuation of Important Safety Information on [page 11](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.

oxervate® 
(cenergermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

Using OXERVATE

This dosing and administration overview will help you understand how to use OXERVATE®



This image is for illustrative purposes only. This is not an exact representation of a patient.

Before you use OXERVATE, tell your doctor about all of your medical conditions, including if you:

- have an infection in your eye. If you get an eye infection **while** using OXERVATE, talk to your doctor right away
- are using any other eye drops
- wear contact lenses
- are pregnant or plan to become pregnant. It is not known if OXERVATE will harm your unborn baby
- are breastfeeding or plan to breastfeed. It is not known if OXERVATE passes into your breast milk. Talk to your doctor about the best way to feed your baby if you use OXERVATE

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Only you and your doctor can decide if OXERVATE is right for you

Important Safety Information (cont'd)

What should I avoid while using OXERVATE?

- Your vision may be blurred for a short time after using OXERVATE. If this happens, wait until your vision clears before you drive or use machines.

Please see continuation of Important Safety Information on [page 12](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

How to use OXERVATE®

- See the complete Instructions for Use within the full Prescribing Information, located in pocket, for detailed instructions about the right way to use OXERVATE
- Use OXERVATE exactly as your doctor tells you
- Use 1 drop of OXERVATE in the affected eye or both eyes if needed, 6 times each day, about 2 hours apart starting in the morning. Continue your treatment for 8 weeks
- If you use any other eye drops, wait **at least 15 minutes before or after** using OXERVATE. This will help to avoid one eye drop diluting the other eye drop
- If you also use an eye ointment or gel or an eye drop that is thick, use OXERVATE **first**, and then wait **at least 15 minutes before** using the other eye ointment, gel, or drops
- If you wear contact lenses in your affected eye or both eyes remove them before using OXERVATE and **wait 15 minutes after** using OXERVATE before reinserting them
- If you miss a dose of OXERVATE, take your next dose at your scheduled time. **Do not** take an extra dose to make up for a missed dose
- **Do not** use other eye medicines without talking to your doctor
- Talk to your doctor first before you stop using OXERVATE

If you have any questions about how to use OXERVATE, ask your doctor or pharmacist



This image is for illustrative purposes only. This is not an exact representation of a patient.

What should I avoid while using OXERVATE?

- Your vision may be blurred for a short time after using OXERVATE. If this happens, **wait until your vision clears before you drive or use machines.**

Important Safety Information (cont'd)

What are the possible side effects of OXERVATE?

- The most common side effects of OXERVATE are eye pain, corneal deposits, feeling of having something in the eye, enlarged blood vessels in the white of the eyes, swelling of the eye, sensitivity to light, increase of tears, and headache.

Please see continuation of Important Safety Information on [page 13](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

Dosing schedule

Use OXERVATE® exactly as your doctor tells you

Start in the morning, or as directed by your doctor.



Use **1 drop** in your affected eye(s)



6 times each day, about 2 hours apart

Continue your treatment for **8 weeks**

When getting ready to apply OXERVATE, remember the **15-minute reminders**

If you:

Make sure to:

Use any other eye drops

Use other eye drops **at least 15 minutes before or after** using OXERVATE

Use any other eye ointment, gel, or thick eye drop

Use OXERVATE **first**, then wait **at least 15 minutes before** using the other eye ointment, gel, or drop

Wear contact lenses in your affected eye(s)

Remove them before using OXERVATE. **Wait 15 minutes after** using before reinserting them


Important Safety Information (cont'd)

What are the possible side effects of OXERVATE? (cont'd)

- These are not all the possible side effects of OXERVATE.
- Tell your doctor if you have any side effects that bother you or do not go away.

Please see continuation of Important Safety Information on [page 16](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)



How to use

Not an actual patient.

Please see continuation of Important Safety Information on [page 16](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

How to use

OXERVATE® is supplied in an insulated pack, in weekly cartons containing 7 multiple-dose vials. This may include either frozen and/or refrigerated gel packs or, in some cases, dry ice. OXERVATE is supplied with an accompanying Delivery System Kit. The Delivery System Kit will have medical devices for withdrawing and using OXERVATE. You will receive both the OXERVATE weekly carton(s) and the Delivery System Kit from the pharmacy.

OXERVATE Weekly Carton contains:



7 multiple-dose vials of OXERVATE
(1 vial per day of the week)

Delivery System Kit contains:




7 vial adapters
(+1 extra)



42 pipettes
(+3 extra)



42 sterile disinfectant wipes
(+3 extra)



1 Dose Recording Card

Use the Dose Recording Card included in your Delivery System Kit to:

- Track your 6 doses per day
- Write down the date you start a new weekly supply
- Note the time you opened a new vial (when you connect the vial adapter to the vial) during the week

Important:

- OXERVATE is for use in the eye.
- **Do not** shake the OXERVATE vial.
- Use OXERVATE with the vial adapters, sterile disinfectant wipes, and pipettes that come with your Delivery System Kit.

Please see continuation of Important Safety Information on [page 16](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information for OXERVATE](#).

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

An overview of applying OXERVATE®

Applying OXERVATE is a multistep process using the materials in the Delivery System Kit and should only be done according to the Instructions for Use within the full Prescribing Information.

Prepare the vial



Important Safety Information (cont'd)

What is the most important information I should know about OXERVATE?

- If you wear contact lenses in your affected eye or both eyes, remove them before using OXERVATE and **wait 15 minutes after** using OXERVATE before reinserting them.
- OXERVATE may cause an eye reaction such as eye pain. Talk to your doctor if you experience a more serious eye reaction.
- If you get an eye infection while using OXERVATE, talk to your doctor right away.

Please see continuation of Important Safety Information on [page 24](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

Fill the pipette



Instill a drop



These are not the complete administration instructions.

Scan the codes below for complete information on how to administer OXERVATE or refer to the complete Instructions for Use found within the Prescribing Information.

Dosing and administration video

Scan to watch:
oxervate.com/watch



Complete Instructions for Use

Scan to read:
oxervate.com/IFU



oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)



Support throughout your journey

Not an actual patient.

Two teams are dedicated to helping you access and afford OXERVATE[®] and answer questions you may have

For more information on how Dompé CONNECT to Care and Accredo can support you through the fulfillment and treatment process, [go to page 20](#).

oxervate[®] 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

Dompé CONNECT to Care

may help you access your OXERVATE® prescription:

When your doctor prescribes OXERVATE, you may also enroll in Dompé CONNECT to Care, a patient support program provided by the manufacturer of OXERVATE to help with:

- Insurance verification
- Financial assistance
- Prior authorizations
- General questions

Speak with a Dompé CONNECT to Care representative:

1-877-422-4412

Mon – Fri

8 AM – 8 PM ET

Accredo Health Group Inc., a specialty pharmacy,* manages deliveries of OXERVATE directly to you:

With OXERVATE, you don't go to the pharmacy every time you need a refill. Instead, Accredo ships OXERVATE to your home every 2 weeks, throughout your 8-week course of treatment.

In addition to coordinating your OXERVATE deliveries, an Accredo Pharmacist can also answer questions about dosing, administration, and what to expect during treatment.

Speak with an Accredo representative:

1-888-454-8860

24 hours a day,

7 days a week

*Unlike retail pharmacies, a specialty pharmacy only provides medications used to treat rare or complex diseases or conditions.

Please see continuation of Important Safety Information on [page 24](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

The 5-step fulfillment process

Now that you're familiar with your support teams, let's look at how they'll support you through your OXERVATE® experience:



When you elect to enroll in Dompé CONNECT to Care, you and your doctor will complete the enrollment form

- Within 24 hours, Dompé CONNECT to Care will call to confirm your insurance coverage, so be sure to have your insurance prescription card ready



Dompé CONNECT to Care calls about payment

- Once your insurance is verified, Dompé CONNECT to Care will call again to discuss financial assistance options



Scan the code to add Dompé CONNECT to Care to your contacts so you recognize the number



Accredo calls to coordinate shipment

- After financial assistance is approved, you must speak with an Accredo representative to provide your delivery address



Scan the code to add Accredo to your contacts so you recognize the number

Please see continuation of Important Safety Information on [page 24](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

Step 4



Accredo delivers OXERVATE® to you

- OXERVATE is shipped directly to patients in an insulated pack. This may include either frozen and/or refrigerated gel packs or, in some cases, dry ice

Step 5



Treatment begins

- You're ready to get started

Storage guidelines:

- Remove the weekly carton(s) of OXERVATE from the insulated pack and store for up to 14 days in a refrigerator in the original carton as soon as you can
- Store the weekly carton(s) at 36 °F to 46 °F (2 °C to 8 °C) **no later than 5 hours** from when you receive the medicine from your pharmacy. **Do not** freeze
- OXERVATE is stored in a freezer at the pharmacy. If you start treatment right away after receiving the weekly carton, you will have to wait until the first vial is thawed. Thaw the vial at room temperature up to 77 °F (25 °C). Thawing the vial could take up to 30 minutes when kept at room temperature
- **Keep all medicines out of the reach of children**

Please see continuation of Important Safety Information on [page 24](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)



Not an actual patient.

oxervate® 
(cenegermin-bkbj ophthalmic
solution) 0.002% (20 mcg/mL)

Important Safety Information

What is the most important information I should know about OXERVATE?

- If you wear contact lenses in your affected eye or both eyes, remove them before using OXERVATE and **wait 15 minutes after** using OXERVATE before reinserting them.
- OXERVATE may cause an eye reaction such as eye pain. Talk to your doctor if you experience a more serious eye reaction.
- If you get an eye infection while using OXERVATE, talk to your doctor right away.

What should I avoid while using OXERVATE?

- Your vision may be blurred for a short time after using OXERVATE. If this happens, **wait until your vision clears before you drive or use machines.**

What are the possible side effects of OXERVATE?


- The most common side effects of OXERVATE are eye pain, corneal deposits, feeling of having something in the eye, enlarged blood vessels in the white of the eyes, swelling of the eye, sensitivity to light, increase of tears, and headache.
- These are not all the possible side effects of OXERVATE.
- Tell your doctor if you have any side effects that bother you or do not go away.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088. You can also contact Dompé U.S. Inc. at 1-833-366-7387.

For more information about OXERVATE talk to your doctor or pharmacist.

Please read the full Prescribing Information, including Patient Information, for OXERVATE.

oxervate® 
(cenegermin-bkbj ophthalmic
solution) 0.002% (20 mcg/mL)



Preparing for treatment

You may have questions when you're prescribed OXERVATE®. Here are sample questions to consider asking your doctor.

Not an actual patient.

Important Safety Information (cont'd)

What is the most important information I should know about OXERVATE?

- If you wear contact lenses in your affected eye or both eyes, remove them before using OXERVATE and **wait 15 minutes after** using OXERVATE before reinserting them.

Please see continuation of Important Safety Information on [page 25](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.

oxervate® 
(cenergermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

Why was I prescribed OXERVATE®?

What is neurotrophic keratitis (NK)?

Where do I get OXERVATE?

Do I have to keep OXERVATE refrigerated at all times?

Important Safety Information (cont'd)

What is the most important information I should know about OXERVATE? (cont'd)

- OXERVATE may cause an eye reaction such as eye pain. Talk to your doctor if you experience a more serious eye reaction.
- If you get an eye infection while using OXERVATE, talk to your doctor right away.

What should I avoid while using OXERVATE?

- Your vision may be blurred for a short time after using OXERVATE. If this happens, **wait until your vision clears before you drive or use machines.**

Please see continuation of Important Safety Information on [page 26](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.



I've never used an eye dropper like this. How does it work?

What are the possible side effects of OXERVATE®?

What kind of eye pain might I experience?

When can I expect to see results?

Important Safety Information (cont'd)

What is the most important information I should know about OXERVATE® (cont'd)

- OXERVATE may cause an eye reaction such as eye pain. Talk to your doctor if you experience a more serious eye reaction.
- If you get an eye infection while using OXERVATE, talk to your doctor right away.

Please see continuation of Important Safety Information on [page 27](#) and complete Important Safety Information on [page 23](#), as well as the [full Prescribing Information](#), including [Patient Information](#) for OXERVATE.

oxervate® 
(cenergermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)

Important details about your OXERVATE® treatment:

For more information, go to [OXERVATE.com](https://www.oxervate.com)



OXERVATE® is a prescription eye drop solution used to treat a condition called neurotrophic keratitis. OXERVATE is safe and effective in children two years of age and older.



What are the possible side effects of OXERVATE?

- The most common side effects of OXERVATE are eye pain, corneal deposits, feeling of having something in the eye, enlarged blood vessels in the white of the eyes, swelling of the eye, sensitivity to light, increase of tears, and headache
- These are not all the possible side effects of OXERVATE
- Tell your doctor if you have any side effects that bother you or do not go away

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.



It's important to follow the full 8-week dosing schedule

- Use 1 drop in the affected eye(s) 6 times a day, about 2 hours apart, for 8 weeks

This brochure does not take the place of talking to your doctor about your medical condition or your treatment.

Important Safety Information

What is the most important information I should know about OXERVATE?

- If you wear contact lenses in your affected eye or both eyes, remove them before using OXERVATE and **wait 15 minutes after** using OXERVATE before reinserting them.
- OXERVATE may cause an eye reaction such as eye pain. Talk to your doctor if you experience a more serious eye reaction.
- If you get an eye infection while using OXERVATE, talk to your doctor right away.

Please see complete Important Safety Information on page 23, as well as the full Prescribing Information, including Patient Information for OXERVATE.



© 2025 Dompé U.S. Inc. All rights reserved.
US-OXE-2000067.2 06/25

oxervate® 
(cenegermin-bkbj ophthalmic solution) 0.002% (20 mcg/mL)